DOI: 10.1111/1365-2745.14360

RESEARCH ARTICLE

Journal of Ecology

Forest growth resistance and resilience to the 2018–2020 drought depend on tree diversity and mycorrhizal type

Lena Sachsenmaier^{1,2} | Florian Schnabel^{1,2,3} | Peter Dietrich⁴ | Nico Eisenhauer^{1,5} | Olga Ferlian^{1,5} | Julius Quosh^{1,5} | Ronny Richter^{1,2} | Christian Wirth^{1,2,6}

¹German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany; ²Systematic Botany and Functional Biodiversity, Leipzig University, Leipzig, Germany; ³Institute of Forest Sciences, University of Freiburg, Freiburg, Germany; ⁴Geobotany and Botanical Garden, Martin-Luther-University Halle-Wittenberg, Halle (Saale), Germany; ⁵Institute of Biology, Leipzig University, Leipzig, Germany and ⁶Max-Planck Institute for Biogeochemistry, Jena, Germany

Correspondence

Lena Sachsenmaier Email: lena.sachsenmaier@idiv.de

Funding information

Deutsche Forschungsgemeinschaft, Grant/Award Number: 319936945/ GRK2324, Ei 862/29-1, Ei 862/31-1 and 202548816; University of Chinese Academy of Sciences

Handling Editor: Justine Karst

Abstract

- The frequency of consecutive drought years is predicted to increase due to climate change. These droughts have strong negative impacts on forest ecosystems. Mixing tree species is proposed to increase the drought resistance and resilience of tree communities. However, this promising diversity effect has not yet been investigated under extreme drought conditions and in the context of complementary mycorrhizal associations and their potential role in improving water uptake.
- 2. Here, we investigate whether tree diversity promotes growth resistance and resilience to extreme drought and whether drought responses are modulated by mycorrhizal associations. We used inventory data (2015–2021) from a young tree diversity experiment in Germany, manipulating tree species richness (1, 2 and 4 species) and mycorrhizal type (communities containing arbuscular mycorrhizal [AM] or ectomycorrhizal [EM] tree species, or both). For all tree communities, we calculated basal area increment in the periods before, during and after drought and used the concepts of resistance and resilience to quantify growth responses to drought.
- 3. We found strong growth declines during the extreme 2018–2020 drought for most tree communities. Contrary to our hypothesis, we did not find that tree species richness per se can buffer the negative impacts of extreme drought on tree growth. However, while for EM communities, drought resistance and resilience decreased with tree species richness, they increased for AM communities and communities comprising both mycorrhizal types. We highlight that among various mixtures of tree species, only those with mixed mycorrhizal types outperformed their respective monocultures during and after drought. Furthermore, under extreme drought, the community tends to segregate into 'winner' and 'loser' tree species in terms of diversity, indicating a possible intensification of competition.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

COLOGICAI

- 4. While we cannot disentangle the underlying mechanisms or clarify the role of mycorrhiza during drought, our findings suggest that mixtures of mycorrhizal types within tree communities could help safeguard forests against increasing drought frequency.
- Synthesis. Drought resistance and resilience of tree communities depend on tree diversity and mycorrhizal association types. Mixing tree species with diverse mycorrhizal types holds promise for forest restoration in the face of climate change.

KEYWORDS

biodiversity-ecosystem functioning, climate change, drought, growth resilience, growth resistance, mycorrhizal associations, tree diversity

1 | INTRODUCTION

Global warming leads to an increased likelihood of severe and consecutive droughts (Hari et al., 2020; Spinoni et al., 2018). Forest ecosystems, in particular, face threats from these droughts, given the long generation time and slow growth of trees compared to other plants (Brodribb et al., 2020). This is of great concern because, in addition to preserving biodiversity conservation and other ecosystem services, forests play a crucial role as carbon sinks, thus contributing to the mitigation of present and future climate change (Anderegg et al., 2020).

In 2018, Northern and Central Europe experienced an extraordinary compound drought, characterized by insufficient precipitation and heatwaves (Zscheischler et al., 2020; Zscheischler & Fischer, 2020). The exceptionally dry soil conditions persisted in 2019 and in vast areas of Central Europe, they lasted even through the year of 2020 (Rakovec et al., 2022). The occurrence of these three consecutive drought years from 2018 to 2020 (hereafter referred to as the '2018-2020 drought') marks an unprecedented drought situation in Central Europe, at least within the last 250 years (Bastos et al., 2021; Hari et al., 2020; Rakovec et al., 2022). Such compound and consecutive droughts are substantially increasing in frequency (Hari et al., 2020; Markonis et al., 2021), and accumulating scientific evidence highlights their negative impacts on ecosystems, especially on forests ecosystems (Bastos et al., 2020, 2021; Forzieri et al., 2021; Gampe et al., 2021). Several studies reported widespread premature leaf senescence in 2018, unprecedented droughtinduced tree mortality across various species throughout the region and reductions in tree growth (Bose et al., 2020; Brun et al., 2020; Buras et al., 2020; Schnabel et al., 2022; Schuldt et al., 2020). Tree stress responses were found to be even more pronounced in 2019 than in 2018, indicating that consecutive and compound drought years represent a novel stressor for forests (Schnabel et al., 2022). Lags in physiological recovery, that is, drought legacy effects, can be caused by hydraulic damage (Anderegg et al., 2018; Kannenberg et al., 2019), carbon depletion or shifts in carbon allocation (e.g. towards rebuilding the canopy, growing roots or reproduction), which manifest in the reduction of radial stem growth.

For the quantification of drought impacts, the concept of resistance and resilience could be applied to forest communities to unravel these different facets of ecosystem stability (Ingrisch & Bahn, 2018; Isbell et al., 2015; Lloret et al., 2011). Although there are various approaches to address components of stability (for an overview, see Ingrisch & Bahn, 2018), we consider resistance here as the ability to persist and maintain functioning during a disturbance quantified as the ratio between tree growth during drought and tree growth during the respective pre-drought period, characterized by 'normal' climate conditions (Lloret et al., 2011). Similarly, resilience is defined as the capacity to reach pre-disturbance performance levels, and is estimated as the ratio between post-drought growth and pre-drought growth (Lloret et al., 2011).

Therefore, evaluating the differences in drought resistance and resilience among tree communities represents a crucial step towards understanding how the impacts of drought on forests could be mitigated by the choice of tree species and the design of climate-smart mixtures (Messier et al., 2022).

Biodiversity is known to stabilize ecosystem productivity over time and is therefore considered a key feature that supports the resistance and resilience of ecosystem functions to droughts (Cardinale et al., 2013; Isbell et al., 2015; Jourdan et al., 2020; Morin et al., 2014). In forest ecosystems, the influence of diversity on drought resistance and resilience is attributed to beneficial interactions among tree species such as resource partitioning (e.g. differential stomatal regulation strategies), facilitation (e.g. active hydraulic redistribution) or selection effects (e.g. competitive dominance of deep-rooted species) (Grossiord, 2020). Experimental evidence from subtropical tree communities suggests that the stabilizing effect of tree species richness is driven by interannual variations in the growth of different tree species, which buffer the community against stress-related growth declines (Schnabel et al., 2021).

However, it remains unknown whether this positive diversity effect persists when communities experience an unprecedented drought episode—such as the 2018-2020 drought. Indeed, there are indications that the positive diversity effects observed under moderate drought stress may shift to negative effects due to competitive species interactions (Haberstroh & Werner, 2022). Evidence on the impact of tree diversity on forest growth during and after drought remains inconsistent with positive, but also neutral, and negative diversity effects on tree responses to drought being

Journal of Ecology

FCOLOGICA

reported (Forrester et al., 2016; Gillerot et al., 2021; Grossiord, 2020; Grossiord et al., 2014; Jucker et al., 2014; Pardos et al., 2021).

One possible factor explaining these inconsistent results may be the type of mycorrhizal association of the tree communities. Mycorrhizal fungi are known to help plants acquire nutrients and water uptake in exchange for photosynthates (Bowles et al., 2018; Lehto & Zwiazek, 2011), as mycorrhizal hyphae reach soil water and nutrients that would be inaccessible to plant roots (Allen, 2007). Therefore, the type of mycorrhizal association could play an important role in the effects of drought on forests. A growing body of research suggests that the type of mycorrhizal association is a key driver for ecosystem-functioning relationships (Deng et al., 2023; Luo et al., 2023; Mao et al., 2023). There are two main groups of mycorrhizal association types that are formed between temperate tree species and fungi: ectomycorrhiza (EM) and arbuscular mycorrhiza (AM), which differ in their morphology, physiology and therefore soil nutrient uptake processes (Phillips et al., 2013; Tedersoo & Bahram, 2019). Ectomycorrhizal fungi develop a mantle of hyphae around the tips of the plant roots through which the exchange of nutrients with their hosts occurs. Arbuscular mycorrhizal fungi are endophytic and exchange nutrients within the inner cortical cells of the plant host fine roots (Peterson & Massicotte, 2004). While AM fungi primarily provide their plant host with access to soil phosphorus in the upper mineral soil layer, EM fungi can mobilize both organic and mineral plant resources and typically thrive in organic soil horizons (Midgley & Phillips, 2014; Phillips et al., 2013; Read & Perez-Moreno, 2003; Rosling et al., 2016; Toju et al., 2016). The differences between mycorrhizal types in relation to the benefit provided to their host trees under drought conditions remain unclear, due to a lack of comparative studies (Allen, 2007; Gehring et al., 2006; Kilpeläinen et al., 2017; Mohan et al., 2014; Querejeta et al., 2009; Teste et al., 2020). Although dual mycorrhization with AM and EM in plant roots seems common (Teste et al., 2020), one of the two mycorrhizal types dominates the association in most temperate tree species (Ferlian et al., 2021; Heklau et al., 2021). Due to the distinct lifestyles and foraging strategies of AM and EM fungi, it can be expected that the presence of both types of association within a plant community could lead to higher resource partitioning among their associated plant hosts (Ferlian et al., 2018; Luo et al., 2018; Teste et al., 2020; Wagg, Jansa, Schmid, et al., 2011; Wagg, Jansa, Stadler, et al., 2011). Particularly during drought stress, when tree communities lack water and nutrient supply, the potential benefit of mycorrhizal type richness, which can offer support by complementary resource access (Ferlian et al., 2018; Teste et al., 2020), may become more pronounced.

However, evidence of the promising role of mycorrhizal associations in drought resistance and resilience of tree communities and especially studies using both types of mycorrhizal association are lacking so far. For the determination of management actions for forests in future climatic conditions, it is crucial to understand the relevance of both tree diversity and below-ground mycorrhizal associations, for forest resistance and resilience to drought (Eisenhauer et al., 2022). Such insights could be best achieved with an experimental approach, which manipulates both factors while controlling for confounding environmental effects (Eisenhauer et al., 2022; Ferlian et al., 2018; Scherer-Lorenzen et al., 2005).

Here, we evaluated the growth resistance and resilience of tree communities varying in tree species richness to the 2018–2020 drought, using inventory data from a tree diversity experiment in Germany (MyDiv) that crosses tree species richness (monocultures, 2-species mixtures, 4-species mixtures) with mycorrhizal association types (AM only, EM only and AM+EM).

Specifically, we tested the following hypotheses:

H1. Growth resistance and resilience to the 2018–2020 drought increase with tree species richness.

H2. Communities that contain tree species of both mycorrhizal association types (AM+EM) exhibit higher growth resistance and resilience to the 2018–2020 drought compared to tree communities with either AM associations or EM associations alone.

H3. The relationship between growth resistance and resilience to the 2018–2020 drought and tree species richness is modulated by the mycorrhizal association type of the tree communities.

2 | MATERIALS AND METHODS

2.1 | Study site and experimental design

This study was carried out in the MyDiv Experiment, a tree diversity experiment located at the Bad Lauchstädt Experimental Research Station of the Helmholtz Centre for Environmental Research-UFZ, close to Halle (Saale) in Germany (51°23' N, 11°53' E; 118 ma.s.l.). The climate at the site is continental summer dry with a mean annual precipitation sum of 484 mm and a mean temperature of 8.8°C (1896-2003). The soil at the site is a Haplic Chernozem developed from loess (Altermann et al., 2005) and was formerly used agriculturally until 2012 and as grassland until the establishment of the experiment in 2015. The study site was divided into two blocks based on the abiotic and biotic parameters measured before planting (Ferlian et al., 2018). The experiment consists of 80 plots, each 11 m × 11 m in size. In each plot, 140 two-year-old tree saplings were planted on a regular grid with a distance of 1 m between individuals. Tree species were selected from a species pool of 10 common temperate deciduous tree species, five identified in the literature as predominantly associated with EM fungi and five predominantly with AM fungi (Table 1). The experiment is designed with a tree species richness gradient ranging from monocultures to 4-species mixture plots (1, 2 and 4 species), crossed with the plot mycorrhizal association type, determined by the selected tree species. This results in Journal of Ecology 🗧

SACHSENMAIER ET AL.

communities that comprise only AM tree species (n = 10 for each diversity level), only EM tree species (n = 10 for each diversity level), or a combination of AM and EM tree species (AM + EM) (n = 10 for diversity levels 2 and 4). Although the mycorrhizal design was established only indirectly by selecting tree species, a 2019 study (Ferlian et al., 2021) provided confirmation that EM and AM trees are predominantly colonized by EM and AM fungi, respectively.

2.2 | Identification of drought period

We define drought as a period with higher water deficits in comparison to normal conditions, that is, long-term means of meteorological parameters (Schwarz et al., 2020). We used the Standardized Precipitation Evapotranspiration Index (SPEI; Vicente-Serrano et al., 2010) and soil moisture patterns for the determination of drought years at our site. With this approach, we could quantify the severity of drought on different time scales based on a commonly used and standardized index, while also considering the importance of local soil conditions for plant-available water (Schwarz et al., 2020).

The SPEI employs the concept of 'climatic water balances', calculated as available water (precipitation) minus atmospheric evaporative demand (reference evapotranspiration) at different time scales. The resulting water balances are standardized by fitting a log-logistic probability distribution, ensuring comparability across space and time (for more details, see Beguería et al., 2014; Vicente-Serrano et al., 2010). SPEI series were calculated with the SPEI package (Beguería et al., 2014) in R (R Core Team, 2023) from monthly precipitation (mm) and potential evapotranspiration (mm; computed following the Penman-Monteith equation, Allen et al., 1998) data derived from the weather station located closest to the study site and with continuous records (DWD Climate Data Center [CDC], Station Leipzig/Halle, ID 2932; Figure S2). The last 40 years (1982-2022) were used as a reference period. With the calculated SPEI values, we classified the years between 1982 and 2022 as normal $(SPEI \ge (-1)| \le (+1))$, particularly dry $(SPEI \le (-1))$ or particularly wet (SPEI \geq (+1)) (Figure S2). The years 2016 and 2017 can be considered normal years, as their SPEI values are between (+1) and (-1). The years 2018, 2019 and 2020 had the lowest SPEI values in the last 40 years, that is, we identified them as 3 years of consecutive severe drought. The year 2021 can be considered a particularly wet year, according to

TABLE 1 Tree species characteristics and performances in the MyDiv experiment, grouped by their dominant mycorrhizal association type.

Tree species	Family	Species code	Leaf symbol	Dominant mycorrhizal type	Mortality (%) (until 2021)	Mean height (m) (in 2021)	Mean growth rate (cm ² /year) (2016–2021)
Acer pseudoplatanus L.	Sapindaceae	Ac	*	AM	4.7	5.6	8.4
Aesculus hippocastanum L.	Sapindaceae	Ae	﴾	AM	2.9	3.2	4.5
Fraxinus excelsior L.	Oleaceae	Fr	*	AM	10.7	4.8	6.7
Prunus avium (L.) L.	Rosaceae	Pr	-	AM	1.4	5.7	9.7
Sorbus aucuparia L.	Rosaceae	So		АМ	13.5	4.5	5.1
Betula pendula Roth	Betulaceae	Be		EM	21.9	6.0	10.9
Carpinus betulus L.	Betulaceae	Ca		EM	1.2	4.6	5.0
Fagus sylvatica L.	Fagaceae	Fa		EM	12.7	2.6	1.8
Quercus petraea (Matt.) Liebl.	Fagaceae	Qu		EM	12.1	2.8	2.1
Tilia plathyphyllos Scop.	Malvaceae	Ti		EM	2.0	4.5	8.0

its SPEI value higher than (+1). On site soil moisture data since 2017 that were collected by loggers directly in the experimental plots confirmed this drought year identification based on the SPEI (Figure S2).

2.3 | Tree growth responses

2.3.1 | Tree measurements

In repeated annual tree inventories from 2015 to 2021, individual tree stem diameters were measured 5 cm above the ground with a diameter tape (basal diameter, *d*0; cm) in all 80 plots of the experiment. To avoid potential edge effects, we only used an area of 6×6 m and 36 tree individuals per plot, resulting in 2880 trees in total.

2.3.2 | Data cleaning and preparation

Measurement errors in tree inventory data with an annual resolution are common due to, for example, inconsistencies in the precise measurement position at the stem, inconsistencies in the selection of the measured main stem for trees with multiple stems or by the breakage and regrowth of a new stem in the same year (Fichtner et al., 2018; Schnabel et al., 2019). We applied a correction procedure for 4.2% of the values in the data set, that is, we predicted the basal diameter of the tree by constructing individual-based allometric models with the use of the following additional variables (a) diameter at breast height in case the tree was higher than 1.3 m, or (b) height in case the tree was lower than 1.3 m (see Supporting Information, Section III for more details).

Before analysis, we excluded tree individuals with incomplete measurement series over 7 consecutive years (2015–2021) (e.g. due to mortality, wind breakage and resprouting in the next year). Additionally, we removed all tree individuals from one plot of the experiment (monoculture of *Betula pendula*) due to its overall high damage and mortality caused by a storm event. This led to the exclusion of 272 tree individuals (9%) from the data set.

With this approach, there is a potential to overlook poorly performing trees, which could either restrict or enhance observed diversity effects. In general, to protect against any bias introduced by mortality, we calculated a mortality variable for each plot and year. This variable reflects the cumulative basal area lost as a percentage of the total basal area on the plot and was incorporated into all models as an additive factor. Importantly, subsequent analysis revealed that mortality did not emerge as a significant factor in any of the models.

2.3.3 | Community productivity

Using d0, we calculated the tree basal area increment (BAI) for each year as follows:

$$\mathsf{BAI}_{\mathsf{year}} = \left(\pi \times \left(d\mathsf{O}_{\mathsf{year}}/2\right)^2\right) - \left(\pi \times \left(d\mathsf{O}_{\mathsf{year}-1}/2\right)^2\right), \tag{1}$$

Journal of Ecology

where dO_{year} is the diameter of the tree in the respective year and dO_{year-1} its diameter in the previous year. With individual BAI, we calculate community growth responses per year as the mean BAI (cm²) of all trees that grow in the core area of one plot (max. 36 individuals) in the respective year.

2.3.4 | Resistance and resilience

Tree community growth responses to drought were expressed as drought resistance and resilience (Lloret et al., 2011). We based the calculation of these indices on the tree community's mean BAI in the period before (2016-2017; 'pre'), during (2018-2020; 'drought') and after (2021; 'post') the drought. It is important to note that the duration of the pre-disturbance reference period markedly influences the resulting resistance value (Schwarz et al., 2020). However, due to the trees being planted in 2015, our options were limited. As recommended by Schwarz et al. (2020), we compared our results for different reference periods (2016-2017 vs. 2017 only), and the results were found to be consistent. Therefore, we stayed with the more robust 2-year reference period of 2016-2017. With 2022 also experiencing a severe drought, we could only assign 2021 as the post-disturbance year. Acknowledging that the time frame of 1 year may not be optimal for expecting full recovery after a 3-year drought, we refer to this index as early resilience. We calculate resistance, the capacity to withstand drought (Lloret et al., 2011) as:

Resistance
$$(R_t) = \frac{BAI_{drought}}{BAI_{pre}}$$
. (2)

A resistance value below 1 indicates reduced growth during drought compared to the pre-drought period, while a value above 1 indicates increased growth in the drought period compared to the pre-drought period. However, given that we examine very young tree individuals in the phase of exponential radial stem growth, we note that resistance values >1 should be expected and a value around 1 represents an actual decline in growth.

We calculated early resilience, the ability to return to predrought conditions (Lloret et al., 2011), as:

(early) Resilience
$$(R_s) = \frac{BAI_{post}}{BAI_{pre}}$$
. (3)

A resilience value of 1 indicates post-drought growth comparable to the pre-drought period and could result from either (a) high drought resistance or (b) low drought resistance coupled with strong recovery.

2.3.5 | Community overyielding and species-specific overyielding

Using the individual BAI values per tree, we calculated the mean growth performance of the monocultures in the pre-drought, drought and post-drought period for the 10 tree species. Based on these values, we

estimated the expected growth of mixed communities under the assumption that there would be no difference between the effect of interspecific and intraspecific interactions (Forrester & Pretzsch, 2015). Since the trees were regularly planted with the same number of individuals per species in the mixtures (i.e. even mixing proportions per species), we used the mean over all monoculture performances per species as the 'expected' community productivity in mixtures.

We quantified community-level over- or underyielding for each period as

community overyielding [%] =
$$\left(\frac{\text{commBAI}_{\text{mix}}}{\text{commBAI}_{\text{expect}}} - 1\right) \times 100, (4)$$

where commBAI_{mix} represents the actual productivity of the community in a mixture of two or four species, and commBAI_{expect} is the expected productivity of this community, calculated as the mean of all monoculture productivities of the species present in the respective community. When overyielding equals 0, the performance of the mixtures aligns with expectations based on the monocultures. Values above 1 represent improved performance in mixtures, indicating a benefit of mixing, while values below 1 indicate that the mixture performs worse than the corresponding monocultures.

Since a study at the same experimental site revealed that observed patterns of complementarity effects in tree growth were species-specific (Dietrich et al., 2022), we decided to gain some more deeper insights into the underlying mechanisms of tree community behaviour during drought. Through a comparison of overyielding values among the different tree species, we aimed to disentangle the potentially contrasting mixing effects of individual species, which contribute to the diversity effect of the community.

To find out if a certain tree species benefits from growing in a mixture, in particular during drought, we calculated the species-level overyielding as

species overyielding [%] =
$$\left(\frac{\text{speciesBAI}_{mix}}{\text{speciesBAI}_{expect}} - 1\right) \times 100$$
, (5)

using the species monoculture performance (as species BAI_{expect}) and the mean productivity of the species in the respective community (as species BAI_{mix}). A value above 1 indicates that a species benefits from growing in a certain mixture, compared to its performance in monoculture.

2.4 | Statistical analysis

To test our hypothesis that species richness and mycorrhizal association type shape tree community drought resistance and (early) resilience, we generated two linear mixed effect models (LMM) for resistance and resilience, respectively. Logarithmic transformations were applied to the response variables, resistance and resilience, to achieve a normal distribution of the residuals. We modelled resistance and resilience in response to the fixed effects of tree species richness (1, 2 and 4; log-transformed), mycorrhizal type (as a factor: AM, AM+EM and EM) and their interaction. Furthermore, we included the mean tree basal area per community across years (tree size) and the accumulated lost plot basal area (mortality) as fixed effects to control for the effects of tree size and mortality and used the experimental block as a random effect.

Since we were not interested in examining the development of community productivity over the years and observed that absolute productivity was affected by individual species with high absolute growth rates (see Figure S6; Table 1), we focused on analysing relative responses. Consequently, we used an LMM to test whether the overyielding of mixed communities depended on drought and mycorrhizal type using overyielding as the response variable, and drought period (pre, drought, post), mycorrhizal type (AM, AM+EM and EM) and their interaction as fixed effects. We also included species richness (as a factor: 2-, and 4-species) and mortality (as accumulated lost plot basal area) as additional fixed effects and controlled for repeated measurements by including the plot ID as a random effect.

We tested for overyielding in individual species and its dependence on period (pre, drought, post), through an LMM predicting species overyielding by interactive fixed effects of species identity (10 tree species), period (pre, drought, post) and tree species richness (2-, 4-species), with mortality (as accumulated lost plot basal area) as additional fixed effect and plot ID as a random factor. Additionally, to test if species differ significantly in their overyielding between different drought periods, we used post hoc pairwise comparisons of drought periods within each species with the *contrast* function of the *emmeans* package (Lenth, 2023), corrected for multiple comparisons via Tukey's Honestly Significant Difference adjustment.

For all models, we visually assessed linearity, independence, homoscedasticity and normality of residuals, and checked for the absence of influential outliers and potential multicollinearity of predictor variables using the *performance* package (Lüdecke et al., 2021).

All analyses were conducted in R (version 4.3.1, R Core Team, 2023) using the packages *Ime4* (Bates et al., 2015) for LMMs, *ImerTest* (Kuznetsova et al., 2017) for model selection via likelihood ratio tests, *emmeans* (Lenth, 2023) to extract model results and perform post hoc comparisons and *ggplot2* for graphics (Wickham, 2016).

3 | RESULTS

3.1 | Strong growth reductions during extreme drought

We found pronounced responses to drought in terms of reduced tree growth during the 2018–2020 drought event. In the pre-drought period of 2016/2017, young trees exhibited an average growth (BAI) of $8.1 \text{ cm}^2 (\pm 2.7 \text{ cm}^2)$ (Table S4). However, over the 3 drought years, tree communities experienced an average growth reduction of 36.8% compared to the pre-drought period (Figure 1). When comparing individual drought years, the most extreme impact was observed in 2018, where tree communities exhibited the lowest average growth rate $(4.8 \text{ cm}^2 \pm 1.5 \text{ cm}^2)$, followed by 2019 $(4.9 \text{ cm}^2 \pm 1.7 \text{ cm}^2)$ and 2020 $(5.7 \text{ cm}^2 \pm 2.2 \text{ cm}^2)$. In the first year after drought, 2021, the

Journal of Ecology

FIGURE 1 Mean annual growth of tree communities per year. Tree basal area increment is shown per plot over the years 2016–2021. The darkness of the grey colour shows the tree species richness level of the tree community. Box plots for each year represent the interquartile range with the median indicated by the line inside the box.

> Vlixture basal area increment relative to monoculture [%]



FIGURE 2 Overyielding of the mixtures compared to monocultures. Shown is the basal area increment of mixed tree communities relative to their expected productivity based on the productivity of the constituent monocultures (black 0-line). The coloured points and error bars (95% CI) show fits of the linear mixed effects model that predict mean overyielding based on the examined period, tree mycorrhizal type and species richness (see Section 2). The zero line represents the expected monoculture yield, that is, values above this line indicate a mixture performance that is better than the respective monoculture, and values below this line indicate a mixture performance that is worse than the respective monoculture. The model explained 14% and 48% of the variation in overyielding through its fixed (marginal R^2) and fixed and random effects (conditional R^2). The different panels show the mycorrhizal community type (AM, arbuscular mycorrhiza; AM + EM, both types; EM, ectomycorrhiza).

average growth rate slightly exceeded with $8.4 \text{ cm}^2 (\pm 2.1 \text{ cm}^2)$ the levels observed in the pre-drought period (Table S4). It is essential to note that young trees undergo a phase of exponential growth (Pretzsch, 2020); hence, rather than expecting constant growth, a notable increase in the growth rate would have been anticipated in 2018 and the subsequent years under ambient weather conditions.

Focusing on net diversity effects, we modelled their overyielding in relation to their respective monocultures across different periods of the drought event and with respect to the different mycorrhizal types of the plots. Our findings reveal that both the drought period and the mycorrhizal type interacted significantly as drivers of community overyielding ($F_{4,114}$ = 3.38, p < 0.05; Table S5). The impact of whether the tree community consists of two or four different tree species on this relationship was hereby not significant. Before drought, most diverse tree communities overyielded (Figure 2). However, during the drought period, the communities generally showed reduced overyielding compared to the pre-drought period, approaching the performance levels of monocultures (indicated by

the horizontal line, Figure 2). In the post-drought period, more communities showed underyielding than before or during the drought.

Examining the different mycorrhizal types of the communities explained some of the shifts in overyielding patterns: In the predrought period, EM communities and communities with both mycorrhizal types, but not AM communities, clearly overyielded. We highlight that during drought, only communities with both mycorrhizal types (AM + E.1M) significantly outperformed their respective monocultures (Figure 2). In the post-drought period, communities with AM trees and those with both mycorrhizal types predominantly exhibit overyielding tendencies, whereas EM mixtures tended to underperform compared to their respective monocultures (Figure 2).

3.2 | Resistance and resilience modulated by species richness and mycorrhizal type

Overall, the tree communities had a drought resistance of 0.76 (±0.5), that is, they showed a decrease in growth (R_t < 1) during the drought compared to the pre-drought period. However, the same communities showed a mean drought resilience of 1.30 (±1.2), that is, they increased their growth after drought compared to the period before drought. For both responses, resistance and resilience, we found a high variability between the different communities (Figure 3). Overall, we found that mycorrhizal association type significantly shaped the relationship of tree species in both the community's drought resistance ($F_{2,70}$ =5.23, p=0.0077, conditional R^2 =0.65) and resilience ($F_{2,70}$ =6.63, p=0.0023, conditional R^2 =0.59) (Figure 3; Table S6). Unlike our hypotheses, the richness of tree species did not

consistently increase the drought resistance and resilience of tree communities (Table S6; Figure S7). Instead, tree species richness increased drought resistance and resilience only for AM communities and communities with both mycorrhizal association types, but decreased drought resistance and resilience for EM communities.

In contrast to our hypothesis (H2), EM communities and not communities with mixed mycorrhizal types showed the highest drought resistance and resilience (Figure S7). Within the group of EM communities, monocultures were the most resistant and resilient. However, focusing on tree species mixtures of the experiment, we found that while the mixed mycorrhizal communities showed an intermediate response between the EM and AM communities in the 2-species mixtures, they surpassed both other groups in the 4-species mixtures (Figure 3). The relationship of drought resistance and resilience with tree species richness and mycorrhizal type was found to be additionally significantly dependent on the mean size of the trees in the community, with slightly higher resistance and resilience in small-sized trees (Table S6; Figure S8).

3.3 | Tree species identity determines the diversity benefit

We found that the benefit of diversity was strongly species-specific and depended on the drought period (Figure 4). Our model revealed that the interaction between species and the drought period significantly predicted species overyielding in mixtures ($F_{18,404} = 6.77$, p < 0.001, conditional $R^2 = 0.84$). The degree to which a species benefited from the presence of other tree species in the community or



FIGURE 3 Tree community's (a) drought resistance and (b) early resilience as a function of tree species richness and mycorrhizal association type. The colours refer to the type of mycorrhizal association of the plot. Solid lines show significant ($p \le 0.05$) linear mixed effects model fits and shaded areas show confidence intervals of 95% certainty. The dashed horizontal line at the intercept of y = 1 as visual support for interpretation: The values on this horizontal line are communities that grew as much during drought than before (resistance, a) or that grew as much after drought than before (resistance, b), values above the line stand for very high resistance and resilience, that is, communities grew even more during drought than before (resistance, a) or more after drought than before (resilience, b).

SACHSENMAIER ET AL.



FIGURE 4 Overyielding of individual species in mixtures compared to monocultures. Each data point shows the basal area increment of a species on a mixture plot relative to its performance in monoculture (black 0-line). Coloured points and error bars (95% Cls) show linear mixed-effects model fits that predict mean overyielding based on the examined period, tree species and tree species richness (see Section 2). The zero line represents the expected monoculture yield, that is, values above this line indicate a species performance in mixture that is better than the respective monoculture, and values below this line indicate a performance that is worse than the respective monoculture. The model explained 65% and 84% of the variation in overyielding through its fixed (marginal R²) and fixed and random effects (conditional R²). The different panels show which species are assigned to which mycorrhizal type. Species: Ac, Acer pseudoplatanus; Ae, Aesculus hippocastanum; Be, Betula pendula; Ca, Carpinus betulus; Fa, Fagus sylvatica; Fr, Fraxinus excelsior; Pr, Prunus avium; Qu, Quercus petraea; So, Sorbus aucuparia; Ti, Tilia platyphyllos.

not was modulated by drought stress. Species such as A. *pseudoplatanus*, *P. avium* or *B. pendula*, which already benefited from diversity under normal climatic conditions, experienced even greater advantages during drought. Conversely, species that did not benefit from diversity under normal climatic conditions, such as *A. hippocastanum*, *F. sylvatica* or *Q. petraea*, tended to be even more negatively affected by diversity during drought conditions (Figure 4). These trends were more pronounced in both directions at the highest level of species richness for most species. Still, the lack of a significant modulation by tree species richness in the model implies that the observed effects remained relatively unaffected by the type of mixture (2- or 4-species mixtures) (Tables S8 and S9).

4 | DISCUSSION

4.1 | Strong growth reductions during extreme drought

We found strong growth declines during the 2018–2020 drought event for most of the investigated tree communities (Figure 1). Other studies that assessed the impact of the 2018–2020 drought on forests have consistently reported signs of tree drought stress, reduced growth and increased tree mortality, both during and after the drought period (Buras et al., 2020; Obladen et al., 2021; Pohl et al., 2023; Schnabel et al., 2022; Schuldt et al., 2020; Senf et al., 2020). Although our reported growth reduction by 36.8% lines up with the finding of an earlier study by Thom et al., 2023, that reported a 41.3% reduction during the 2018–2020 drought, it should be kept in mind that our reductions are biologically even higher, because we expect an ontogenetic increase (Pretzsch, 2020). Although a generally favourable nutrient supply can alleviate drought impacts on trees (Schmied et al., 2023), our initial anticipation of less pronounced growth reductions, based on the consideration of the soil characteristics at the site (Altermann et al., 2005)–particularly its high fertility and favourable water relations (see Section 2.1)–did not align with the observed outcomes.

Journal of Ecology

In terms of meteorological severity, out of the three drought years, 2018 stood out as the most extreme, followed by 2019 and 2020 (Hari et al., 2020; Rakovec et al., 2022) (Figures S1 and S3). The growth responses of the tree communities at our site appeared to closely mirror the meteorological drought patterns, with the lowest growth rate in 2018, followed by 2019 and 2020. However, contrary findings from other studies point to 2019 as the year with the strongest growth reduction (Pohl et al., 2023; Salomón

Journal of Ecology 📃

et al., 2022; Schnabel et al., 2022), attributed to the legacy effects of the 2018 drought (Anderegg et al., 2015; Kannenberg et al., 2020; Schnabel et al., 2022). Such drought legacies might be caused by diminished reserves of non-structural carbohydrates and altered carbon allocation, favouring the canopy and root system instead of radial stem growth (Brunner et al., 2015; Hartmann & Trumbore, 2016; Kannenberg et al., 2019). However, compared to most other studies, we studied young tree communities (planted in 2015; 5-6 years old when the drought hit), and the growth reduction we observed in 2019 may be interpreted as a strong reduction relative to the exponential growth trajectory expected for young trees (Pretzsch, 2020). Furthermore, the less developed root system of young trees may have increased their sensitivity and led to earlier drought stress compared to trees in a mature stage, which may explain the pronounced reduction observed in 2018 (Franceschini & Schneider, 2014).

The year following drought–2021–was a year with exceptionally high water supply (Figures S1 and S2). Nevertheless, we observed that tree growth remained lower in 2021 than in 2017–the year immediately preceding the onset of drought. Since the analysed trees experienced 3 consecutive years of extreme drought, it is not surprising that they did not recover completely after a single year– whether this was despite, due to or independently unaffected by their young age, we cannot clarify with our data.

4.2 | Tree diversity did not increase drought resistance and resilience per se

In contrast to our hypothesis, we did not find that tree species richness per se can buffer the negative impacts of extreme drought in 2018-2020. The drought resistance and resilience of the investigated tree communities did not increase consistently with increasing tree species richness, but depended on mycorrhizal association type. Previous studies also provided mixed results for the effect of tree diversity on tree responses to drought (Dănescu et al., 2018; Gillerot et al., 2021; Grossiord, 2020; Jourdan et al., 2020; Pardos et al., 2021). These contrasting results regarding the role of tree diversity during drought may be explained by different drought tolerance strategies of the admixed species (Schnabel et al., 2024) and the intensity of the examined drought event. Under severe drought conditions, diversity effects on tree responses to drought may shift from positive to negative (Haberstroh & Werner, 2022), which is also supported by recent results from another tree diversity experiment in Germany regarding the 2018 drought (Kreinitz experiment, Shovon et al., 2024). Indeed, our findings did not support our expectation that mixed-species tree communities would generally benefit from the use of complementary resources, particularly water, leading to a reduction in drought stress and a decrease in growth compared to monocultures. One possible explanation could be that during extreme water scarcity-such as during the 2018-2020 droughteven trees with complementary resource use strategies compete for water resources (Haberstroh & Werner, 2022). A potentially

enhanced competition level in mixtures during extreme drought is also supported by the species-specific analysis we present below. Moreover, it is important to consider that the higher productivity of mixed communities-which was also found for the mixtures of this study (Table S2; Dietrich et al., 2022)-may increase the demand for water (Ammer, 2019). Mixtures could become more vulnerable to drought, unless mitigated by improved water supply via facilitated uptake (Forrester, 2015). Under extremely dry conditions, such as during the 2018-2020 drought, facilitation mechanisms, such as below-ground niche differentiation, may no longer have been able to maintain tree water supply due to excessively dried soils. In these situations, the potentially advantageous larger root system of larger trees (Hui et al., 2014) within a mixture, would not confer benefits either. Our results reveal a similar response pattern for drought resistance as for early drought resilience (Figure 3). Although we cannot confirm any positive effect of tree diversity on resilience (Anderegg et al., 2018), our ability to draw comprehensive conclusions about drought resilience is limited by the fact that our investigation only spans 1 year following the three consecutive drought years.

While various studies emphasize the influence of tree species composition on drought responses in mixed versus monospecific stands (Dănescu et al., 2018; Gillerot et al., 2021; Jourdan et al., 2020; Pardos et al., 2021), the distinctive strength of our study emerges from being among the first to examine the impacts of the 2018-2020 drought under the controlled conditions of a planted tree diversity experiment with various combinations of tree species exposed to the same abiotic conditions. Capitalizing on this set-up, we revealed that the observed drought effects were not only driven by tree species richness but rather depended on the mycorrhizal association type of the examined tree communities.

4.3 | Mycorrhizal types modulate drought responses

In line with our expectations, we found different responses to drought in AM and EM tree communities, but with higher resistance and resilience for EM tree communities. There is evidence that the mycorrhizal association types differ in their nutrient economy: AM fungi rely on inorganic nutrient resources, and EM fungi have the ability to decompose organic matter (Averill et al., 2019; Deng et al., 2023; Liese et al., 2018; Phillips et al., 2013; Tedersoo & Bahram, 2019; Zhang et al., 2018). However, given that the study site has relatively high soil nutrient availability due to its former agricultural use (Ferlian et al., 2018), the differences between AM and EM in the acquisition of nitrogen and phosphorus, may not be decisive in this context. In particular, during drought, it might be more advantageous for tree communities to have an enhanced water supply through their fungal partners than an improved nutrient supply. Both mycorrhizal fungi have mechanisms to maintain host vitality under drought, such as the induction of host aquaporin expression, regulating water uptake (Allen, 2007; Lehto & Zwiazek, 2011; Mohan et al., 2014; Tedersoo & Bahram, 2019; Xu & Zwiazek, 2020). The

question of whether EM or AM associations offer greater drought resistance and resilience to their host trees remains uncertain, given the contrasting results reported so far and the lack of studies that compare both mycorrhizal types (Mohan et al., 2014; Querejeta et al., 2009). Some studies would support the benefits of AM associations under drought, such as AM hyphae being able to endure highly negative water potentials and having a greater plasticity of hyphal production, which might support the existence of AM host plants in extremely water-limited systems (Querejeta et al., 2007; Tedersoo & Bahram, 2019; Vargas et al., 2010). A few studies on dual-mycorrhizal tree species revealed that dry soil favoured the colonization dominance by AM fungi (Gehring et al., 2006; Kilpeläinen et al., 2017; Querejeta et al., 2009). However, our results show that tree communities with EM associations are more resistant and resilient at our site. We propose the following speculative explanations for this response: EM fungi are expected to transport soil water more efficiently due to their greater mycelium biomass and their ability to build vessel-like rhizomorphs (Allen, 2007). Since EM fungi form a Hartig net of hyphae surrounding the root cortex cells and a hyphal sheath, covering the root tips (Freschet et al., 2021), they offer their host superior physical protection against soil-borne pathogens compared to AM fungi (Tedersoo & Bahram, 2019). This enhanced protection might be particularly advantageous during periods of stress, such as drought. Accordingly, it was observed that limited soil water supply reduced stem biomass production stronger for AM than for EM trees in a mesocosm drought experiment (Liese et al., 2018). Furthermore, the drought treatment only reduced fine root biomass and mycorrhizal colonization rates in AM trees, not in EM trees. On the broader scale of different forest communities across the United States. Luo et al. (2023) found EM-dominated forest communities to be more productive than mixed mycorrhizal communities or AM-dominated ones only in those ecoregions that exhibit low mean annual precipitation, while mixed mycorrhizal communities were generally most productive across all ecoregions. Even though the mechanisms remain unclear, these findings on EM-dominated forests support ours, as the EM tree communities showed the highest drought resistance and resilience at our site. Still, while the evidence of the importance of mycorrhizal types for forest ecosystem functioning relationships is accumulating (Deng et al., 2023; Dietrich et al., 2022; Luo et al., 2023; Mao et al., 2023), we need more knowledge on how these relationships are influenced by drought.

Interestingly, within the EM tree communities in our study, monocultures demonstrated greater resistance and resilience than mixtures, while the opposite trend was observed for AM tree communities, that is, higher resistance and resilience were associated with higher tree species diversity. This could be explained by potentially lower root protection by AM fungi colonization compared to EM fungi colonization, resulting in AM trees suffering more from the accumulation of antagonists near conspecifics compared to EM trees (Bennett et al., 2017; Jiang et al., 2020).

Although some EM monocultures proved to be very resistant and resilient, the large differences between EM species (Figure 3, Journal of Ecology

confidence intervals) and their variability in growth performance across periods (Figures 2 and 4) discourage the recommendation of EM monocultures. Instead, our findings demonstrate the beneficial outcomes of mixing mycorrhizal types. While studies comparing tree communities of both mycorrhizal types are scarce, research on dual-mycorrhizal tree species suggests that these may adjust the abundance of each type in response to various factors, including soil moisture (Teste et al., 2020). At our experimental site, trees consistently preferred one mycorrhizal type but also exhibited mycorrhization of the opposite type, with rates varying throughout the season (Ferlian et al., 2021; Heklau et al., 2021, 2023). Mycorrhizal spillover effects from neighbouring host tree species may affect EM and AM synchrony (Dickie et al., 2001), altering mycorrhization rates and fungal composition at the community level (Heklau et al., 2023), which could potentially lead to the observed positive effects of mycorrhizal type mixing.

In previous studies at the same site, biomass production and stand structural complexity were measured as responses to the mixing of mycorrhizal types. These studies did not find a general beneficial effect of mixing mycorrhizal types; instead, they observed a tendency for an additive effect of EM and AM trees on biomass production (Dietrich et al., 2022; Ferlian et al., 2018) or stand structural complexity (Ray et al., 2023). However, our study revealed a strong positive diversity effect only between mixed mycorrhizal communities, indicating enhanced complementarity under drought conditions (Baert et al., 2018). Mixed mycorrhizal communities showed the highest resistance and resilience within 4-species communities (Figure 3) and outperformed monocultures before, during and after drought (Figure 2), suggesting a novel drought mitigation effect of diversity. The variability in mycorrhizal-type modulation across different periods might explain why studies like Ray et al. (2023) found no consistent mycorrhizal effect on productivity throughout their study at the same site.

While there is not yet much literature to support our findings, recent studies indicate that mixed mycorrhizal strategies can promote ecosystem functioning. For instance, Luo et al. (2023) reported that communities with mixed mycorrhizal strategies were more productive than those dominated by either EM or AM tree species. However, our study is the first to demonstrate the positive effects of mixed mycorrhizal strategies on drought responses. If confirmed in subsequent studies, this positive effect of mycorrhizal type mixtures could offer a novel drought mitigation strategy in forests, complementing the positive effect of tree diversity reported in earlier studies (e.g. Fichtner et al., 2020; Schnabel et al., 2019).

4.4 | Tree species' benefits and disadvantages of diversity get stronger during drought

Our results revealed that the effects of diversity on productivity were highly species-specific. For instance, species such as A. *pseudo-platanus*, P. *avium* and B. *pendula* clearly benefited when growing in a mixture compared to their monocultures. On the other hand, species such as A. *hippocastanum*, F. sylvatica and Q. *petraea* thrived more

in monocultures than in mixtures. Species that benefitted from diversity in terms of overyielding grew even better, while species that underyielded in mixtures grew even less during drought than before or after drought. These trends indicate that drought intensified the competitive differences among dominant and subdominant tree species. Our results suggest enhanced competitive dynamics during extreme drought, supporting the assumption that positive effects of diversity may turn negative beyond a threshold of drought stress (Baert et al., 2018; Haberstroh & Werner, 2022; Shovon et al., 2024) and emphasizing the need to consider drought intensity when discussing biotic interactions. Studies on drought responses of single tree species depending on the diversity of the community or neighbourhood are still rare. Some frequently studied species, such as F. sylvatica, were found to benefit from diversity in terms of growth during drought for adult forest trees (Mölder & Leuschner, 2014; Vannoppen et al., 2020), which contrasts with our findings. But this advantage does not persist for F. sylvatica when mixed with conifers (Leuschner, 2020; Thurm et al., 2016), leading to the conclusion that diversity benefits are largely dependent on neighbour identity and neighbour size (Leuschner, 2020). Others state that tree growth responses are contrasting depending on the intensity of drought and the tree species (Bottero et al., 2021). A recent study on diversity effects during the 2018 drought reported that complementarity effects disappeared in species-rich communities, whereas they were still evident for certain combinations of two tree species (Shovon et al., 2024). In general, it seems to be evident that the influence of competition on tree growth responses during drought does not occur in an unidirectional and universal way for all species (Castagneri et al., 2022; Gillerot et al., 2021). Our results show that highly productive pioneer species, such as B. pendula, P. avium or A. pseudoplatanus, suffered during drought in general (Figure S9); however, they benefitted more from growing in the mixture than the slow-growing species, such as F. sylvatica or Q. petraea. This observation is consistent with other experimental studies that reported that tree species richness particularly supported the most drought-vulnerable species in a community, characterized by acquisitive and water-spending functional traits (Fichtner et al., 2020; Schnabel et al., 2024).

It is important to note that the species-specific growth strategies cannot be clearly attributed to the mycorrhizal type, since growth strategies of, for example, EM tree species were found to cover a broad spectrum from the highest productivity across all species (B. pendula) to the slowest growth rate across all species (F. sylvatica), instead of a uniform strategy in all EM species (Dietrich et al., 2022). Despite efforts to minimize variations in functional traits other than mycorrhizal type in the design of the experiment (Ferlian et al., 2018), we cannot dismiss the possibility that the observed patterns are partly linked to the dynamics of stand development. Furthermore, we assume that, along with species-specific growth strategies, trait-based mechanisms-particularly those regulating water use-may explain the responses we observed. Since water regulation strategies are complex, a perspective of the whole plant (Hartmann et al., 2021) and therefore multiple (hydraulic) traits should be used in future studies to shed light on the drivers

of drought resistance and resilience in addition to the mycorrhizal strategies studied here. Here, we only examined drought stress in terms of tree growth and disregarded other effects such as health deterioration or even mortality. Although existing tree mortality in the experimental plots (Table S2) did not affect our results, the low growth resilience of single communities may be an indicator of future mortality (DeSoto et al., 2020). To gain a more complete image of the 2018–2020 drought consequences, future studies should incorporate additional drought stress indicators, such as carbon isotope ratios (Cherubini et al., 2021; Jucker et al., 2017). Moreover, a thorough examination at the neighbourhood level, where tree-tree interactions occur (Trogisch et al., 2021) may provide a more comprehensive explanation of complementarity and competitive species interactions in the context of drought.

5 | CONCLUSION

Our results showed that it is not tree species diversity per se that modulates drought responses, but it shapes, in interaction with the mycorrhizal types and their diversity, the drought resistance and resilience. Some EM monocultures exhibited the highest levels of resistance and resilience among the studied communities. However, tree communities composed of a single species have other well-known limitations, such as higher variability in growth performances (see Figure 2; Figure S6; Schnabel et al., 2021), and sapling survival rates (Blondeel et al., 2024) relative to mixtures as well as increased susceptibility to specialist pests, pathogens and storms (Jactel et al., 2017; Messier et al., 2022). These considerations undermine the suitability of EM monocultures for fostering resistance and resilience particularly when taking a long-term perspective and when considering multiple stress and disturbance factors. Our findings highlight that among various tree species mixtures, only those with mixed mycorrhizal types consistently exhibited overyielding during the extreme 2018-2020 drought. Even though we cannot elucidate the mechanisms behind the benefit of mixed mycorrhizal types during drought in this study, important consequences can still be drawn from our observations. Our results highlight the potential of mixtures comprising tree species with different mycorrhizal types for effective forest restoration strategies, particularly in the face of an increasing frequency of extreme drought events such as the 2018-2020 drought.

Additionally, we found that the drought-mitigating effect of diversity is most pronounced for fast-growing species, which overall suffered the most from drought, as indicated by their comparable low drought resistance. We observed that drought intensified competitive differences among tree species, resulting in winners and losers under these harsh environmental conditions. Nevertheless, our comprehension of the processes and consequences of drought on interactions within tree species mixtures is still in its early stages. Additional experimental evidence is required to predict the vulnerability of trees in the face of climate extremes. We emphasize the need for: (i) a thorough examination at the neighbourhood level, (ii) detailed information Although the transferability of our results to mature forests is limited, our emphasis on young tree plantations remains particularly pivotal in the context of ongoing reforestation initiatives. Our study is among pioneering efforts to examine the severe impacts of the 2018–2020 drought in an experimental set-up, where tree species richness and mycorrhizal type were manipulated. Capitalizing on this set-up, we could directly compare the influence of mycorrhizal types on the tree communities' drought responses under similar abiotic conditions. Though the topic of mycorrhiza during drought requires further investigations, our findings already imply that a mixture of mycorrhizal types within tree communities may be a promising strategy for safeguarding forests against increasingly frequent severe drought events under climate change.

AUTHOR CONTRIBUTIONS

Christian Wirth, Florian Schnabel and Lena Sachsenmaier conceived the ideas and developed the concept of this study. Nico Eisenhauer and Olga Ferlian designed and established the experiment. Julius Quosh collected the data. Lena Sachsenmaier and Peter Dietrich cleaned the data. Lena Sachsenmaier analysed the data with support from Florian Schnabel, Ronny Richter and Christian Wirth. Lena Sachsenmaier led the writing of the manuscript with support from Florian Schnabel and Christian Wirth. All authors contributed critically to the drafts and gave final approval for publication.

ACKNOWLEDGEMENTS

Thisstudy wassupported by the International Research Training Group TreeDì jointly funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)–319936945/GRK2324 and the University of Chinese Academy of Science (UCAS). N.E. gratefully acknowledges the support of iDiv, which is funded by the German Research Foundation (DFG-FZT 118, 202548816), as well as by the DFG (Ei 862/29-1 and Ei 862/31-1).

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

PEER REVIEW

The peer review history for this article is available at https://www. webofscience.com/api/gateway/wos/peer-review/10.1111/1365-2745.14360.

DATA AVAILABILITY STATEMENT

The data used and analysed during this study are available in the MyDiv database: https://doi.org/10.25829/BPTQ-Z678 (Sachsenmaier et al., 2024).

ORCID

Lena Sachsenmaier b https://orcid.org/0000-0002-0594-3927 Florian Schnabel b https://orcid.org/0000-0001-8452-4001 Peter Dietrich b https://orcid.org/0000-0002-7742-6064 Nico Eisenhauer b https://orcid.org/0000-0002-0371-6720 Olga Ferlian b https://orcid.org/0000-0002-2536-7592 Ronny Richter b https://orcid.org/0000-0002-8728-7918 Christian Wirth b https://orcid.org/0000-0003-2604-8056

REFERENCES

- Allen, M. F. (2007). Mycorrhizal fungi: Highways for water and nutrients in arid soils. Vadose Zone Journal, 6(2), 291–297. https://doi.org/10. 2136/vzj2006.0068
- Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). Crop evapotranspiration-guidelines for computing crop water requirements-FAO irrigation and drainage paper 56 (Vol. 300). Food and Agriculture Organization of the United.
- Altermann, M., Rinklebe, J., Merbach, I., Körschens, M., Langer, U., & Hofmann, B. (2005). Chernozem–Soil of the year 2005. Journal of Plant Nutrition and Soil Science, 168(6), 725–740. https://doi.org/10. 1002/jpln.200521814
- Ammer, C. (2019). Diversity and forest productivity in a changing climate. New Phytologist, 221(1), 50–66. https://doi.org/10.1111/nph.15263
- Anderegg, W. R. L., Konings, A. G., Trugman, A. T., Yu, K., Bowling, D. R., Gabbitas, R., Karp, D. S., Pacala, S., Sperry, J. S., Sulman, B. N., & Zenes, N. (2018). Hydraulic diversity of forests regulates ecosystem resilience during drought. *Nature*, *561*(7724), 538–541. https:// doi.org/10.1038/s41586-018-0539-7
- Anderegg, W. R. L., Schwalm, C., Biondi, F., Camarero, J. J., Koch, G., Litvak, M., Ogle, K., Shaw, J. D., Shevliakova, E., Williams, A. P., Wolf, A., Ziaco, E., & Pacala, S. (2015). Pervasive drought legacies in forest ecosystems and their implications for carbon cycle models. *Science*, 349(6247), 528–532. https://doi.org/10.1126/science. aab1833
- Anderegg, W. R. L., Trugman, A. T., Badgley, G., Anderson, C. M., Bartuska, A., Ciais, P., Cullenward, D., Field, C. B., Freeman, J., Goetz, S. J., Hicke, J. A., Huntzinger, D., Jackson, R. B., Nickerson, J., Pacala, S., & Randerson, J. T. (2020). Climate-driven risks to the climate mitigation potential of forests. *Science*, *368*(6497), eaaz7005. https:// doi.org/10.1126/science.aaz7005
- Averill, C., Bhatnagar, J. M., Dietze, M. C., Pearse, W. D., & Kivlin, S. N. (2019). Global imprint of mycorrhizal fungi on whole-plant nutrient economics. Proceedings of the National Academy of Sciences of the United States of America, 116(46), 23163–23168. https://doi.org/10. 1073/pnas.1906655116
- Baert, J. M., Eisenhauer, N., Janssen, C. R., & De Laender, F. (2018). Biodiversity effects on ecosystem functioning respond unimodally to environmental stress. *Ecology Letters*, 21(8), 1191–1199. https:// doi.org/10.1111/ele.13088
- Bastos, A., Ciais, P., Friedlingstein, P., Sitch, S., Pongratz, J., Fan, L., Wigneron, J. P., Weber, U., Reichstein, M., Fu, Z., Anthoni, P., Arneth, A., Haverd, V., Jain, A. K., Joetzjer, E., Knauer, J., Lienert, S., Loughran, T., McGuire, P. C., ... Zaehle, S. (2020). Direct and seasonal legacy effects of the 2018 heat wave and drought on European ecosystem productivity. *Science Advances*, 6(24), eaba2724. https:// doi.org/10.1126/sciadv.aba2724
- Bastos, A., Orth, R., Reichstein, M., Ciais, P., Viovy, N., Zaehle, S., Anthoni, P., Arneth, A., Gentine, P., Joetzjer, E., Lienert, S., Loughran, T., McGuire, P. C., Sungmin, O., Pongratz, J., & Sitch, S. (2021). Vulnerability of European ecosystems to two compound dry and hot summers in 2018 and 2019. *Earth System Dynamics*, 12(4), 1015–1035. https://doi.org/10.5194/esd-12-1015-2021

Journal of Ecology

- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using Ime4. *Journal of Statistical Software*, 67(1), 1–48. https://doi.org/10.18637/jss.v067.i01
- Beguería, S., Vicente-Serrano, S. M., Reig, F., & Latorre, B. (2014). Standardized precipitation evapotranspiration index (SPEI) revisited: Parameter fitting, evapotranspiration models, tools, datasets and drought monitoring. *International Journal of Climatology*, 34(10), 3001–3023. https://doi.org/10.1002/joc.3887
- Bennett, J. A., Maherali, H., Reinhart, K. O., Lekberg, Y., Hart, M. M., & Klironomos, J. (2017). Plant-soil feedbacks and mycorrhizal type influence temperate forest population dynamics. *Science*, 355(6321), 181–184. https://doi.org/10.1126/science.aai8212
- Blondeel, H., Guillemot, J., Martin-StPaul, N., Druel, A., Bilodeau-Gauthier, S., Bauhus, J., Grossiord, C., Hector, A., Jactel, H., Jensen, J., Messier, C., Muys, B., Serrano-León, H., Auge, H., Barsoum, N., Birhane, E., Bruelheide, H., Cavender-Bares, J., Chu, C., ... Baeten, L. (2024). Tree diversity reduces variability in sapling survival under drought. *Journal of Ecology*, 112(5), 1164–1180. https://doi.org/10. 1111/1365-2745.14294
- Bose, A. K., Gessler, A., Bolte, A., Bottero, A., Buras, A., Cailleret, M., Camarero, J. J., Haeni, M., Hereş, A., Hevia, A., Lévesque, M., Linares, J. C., Martinez-Vilalta, J., Matías, L., Menzel, A., Sánchez-Salguero, R., Saurer, M., Vennetier, M., Ziche, D., & Rigling, A. (2020). Growth and resilience responses of Scots pine to extreme droughts across Europe depend on predrought growth conditions. *Global Change Biology*, 26(8), 4521–4537. https://doi.org/10.1111/gcb.15153
- Bottero, A., Forrester, D. I., Cailleret, M., Kohnle, U., Gessler, A., Michel, D., Bose, A. K., Bauhus, J., Bugmann, H., Cuntz, M., Gillerot, L., Hanewinkel, M., Lévesque, M., Ryder, J., Sainte-Marie, J., Schwarz, J., Yousefpour, R., Zamora-Pereira, J. C., & Rigling, A. (2021). Growth resistance and resilience of mixed silver fir and Norway spruce forests in Central Europe: Contrasting responses to mild and severe droughts. *Global Change Biology*, *27*(18), 4403–4419. https://doi.org/10.1111/gcb.15737
- Bowles, T. M., Jackson, L. E., & Cavagnaro, T. R. (2018). Mycorrhizal fungi enhance plant nutrient acquisition and modulate nitrogen loss with variable water regimes. *Global Change Biology*, 24(1), e171–e182. https://doi.org/10.1111/gcb.13884
- Brodribb, T. J., Powers, J., Cochard, H., & Choat, B. (2020). Hanging by a thread? Forests and drought. *Science*, 368(6488), 261–266. https:// doi.org/10.1126/science.aat7631
- Brun, P., Psomas, A., Ginzler, C., Thuiller, W., Zappa, M., & Zimmermann, N. E. (2020). Large-scale early-wilting response of Central European forests to the 2018 extreme drought. *Global Change Biology*, 26(12), 7021–7035. https://doi.org/10.1111/gcb.15360
- Brunner, I., Herzog, C., Dawes, M. A., Arend, M., & Sperisen, C. (2015). How tree roots respond to drought. Frontiers in Plant Science, 6, 1–16. https://doi.org/10.3389/fpls.2015.00547
- Buras, A., Rammig, A., & Zang, S. C. (2020). Quantifying impacts of the 2018 drought on European ecosystems in comparison to 2003. *Biogeosciences*, 17(6), 1655–1672. https://doi.org/10.5194/ bg-17-1655-2020
- Cardinale, B. J., Gross, K., Fritschie, K., Flombaum, P., Fox, J. W., Rixen, C., Van Ruijven, J., Reich, P. B., Scherer-Lorenzen, M., & Wilsey, B. J. (2013). Biodiversity simultaneously enhances the production and stability of community biomass, but the effects are independent. *Ecology*, 94(8), 1697–1707. https://doi.org/10.1890/12-1334.1
- Castagneri, D., Vacchiano, G., Hacket-Pain, A., DeRose, R. J., Klein, T., & Bottero, A. (2022). Meta-analysis reveals different competition effects on tree growth resistance and resilience to drought. *Ecosystems*, 25(1), 30–43. https://doi.org/10.1007/s10021-021-00638-4
- Cherubini, P., Battipaglia, G., & Innes, J. L. (2021). Tree vitality and Forest health: Can tree-ring stable isotopes Be used as indicators? *Current Forestry Reports*, 7(2), 69–80. https://doi.org/10.1007/s40725-021-00137-8

- Dănescu, A., Kohnle, U., Bauhus, J., Sohn, J., & Albrecht, A. T. (2018). Stability of tree increment in relation to episodic drought in uneven-structured, mixed stands in southwestern Germany. Forest Ecology and Management, 415-416, 148-159. https://doi.org/10. 1016/j.foreco.2018.02.030
- Deng, M., Hu, S., Guo, L., Jiang, L., Huang, Y., Schmid, B., Liu, C., Chang, P., Li, S., Liu, X., Ma, K., & Liu, L. (2023). Tree mycorrhizal association types control biodiversity-productivity relationship in a subtropical forest. *Science Advances*, 9(3), 1–12. https://doi.org/10.1126/ sciady.add4468
- DeSoto, L., Cailleret, M., Sterck, F., Jansen, S., Kramer, K., Robert, E. M. R., Aakala, T., Amoroso, M. M., Bigler, C., Camarero, J. J., Čufar, K., Gea-Izquierdo, G., Gillner, S., Haavik, L. J., Hereş, A. M., Kane, J. M., Kharuk, V. I., Kitzberger, T., Klein, T., ... Martínez-Vilalta, J. (2020). Low growth resilience to drought is related to future mortality risk in trees. *Nature Communications*, 11(1), 1–9. https://doi.org/10. 1038/s41467-020-14300-5
- Dickie, I. A., Koide, R. T., & Fayish, A. C. (2001). Vesicular-arbuscular mycorrhizal infection of Quercus rubra seedlings. *New Phytologist*, 151(1), 257-264. https://doi.org/10.1046/j.1469-8137.2001.00148.x
- Dietrich, P., Ferlian, O., Huang, Y., Luo, S., Quosh, J., & Eisenhauer, N. (2022). Tree diversity effects on productivity depend on mycorrhizae and life strategies in a temperate forest experiment. *Ecology*, 104, e3896. https://doi.org/10.1002/ecy.3896
- Eisenhauer, N., Bonfante, P., Buscot, F., Cesarz, S., Guerra, C., Heintz-Buschart, A., Hines, J., Patoine, G., Rillig, M., Schmid, B., Verheyen, K., Wirth, C., & Ferlian, O. (2022). Biotic interactions as mediators of context-dependent biodiversity-ecosystem functioning relationships. *Research Ideas & Outcomes*, *8*, e85873. https://doi.org/ 10.3897/rio.8.e85873
- Ferlian, O., Cesarz, S., Craven, D., Hines, J., Barry, K. E., Bruelheide, H., Buscot, F., Haider, S., Heklau, H., Herrmann, S., Kühn, P., Pruschitzki, U., Schädler, M., Wagg, C., Weigelt, A., Wubet, T., & Eisenhauer, N. (2018). Mycorrhiza in tree diversity-ecosystem function relationships: Conceptual framework and experimental implementation. *Ecosphere*, 9(5), e02226. https://doi.org/10. 1002/ecs2.2226
- Ferlian, O., Goldmann, K., Eisenhauer, N., Tarkka, M. T., Buscot, F., & Heintz-Buschart, A. (2021). Distinct effects of host and neighbour tree identity on arbuscular and ectomycorrhizal fungi along a tree diversity gradient. *ISME Communications*, 1(1), 1–10. https://doi. org/10.1038/s43705-021-00042-y
- Fichtner, A., Härdtle, W., Bruelheide, H., Kunz, M., Li, Y., & Von Oheimb, G. (2018). Neighbourhood interactions drive overyielding in mixedspecies tree communities. *Nature Communications*, 9(1), 1144. https://doi.org/10.1038/s41467-018-03529-w
- Fichtner, A., Schnabel, F., Bruelheide, H., Kunz, M., Mausolf, K., Schuldt, A., Härdtle, W., & Oheimb, G. (2020). Neighbourhood diversity mitigates drought impacts on tree growth. *Journal of Ecology*, 108(3), 865–875. https://doi.org/10.1111/1365-2745.13353
- Forrester, D. I. (2015). Transpiration and water-use efficiency in mixedspecies forests versus monocultures: Effects of tree size, stand density and season. *Tree Physiology*, 35(3), 289–304. https://doi. org/10.1093/treephys/tpv011
- Forrester, D. I., Bonal, D., Dawud, S., Gessler, A., Granier, A., Pollastrini, M., & Grossiord, C. (2016). Drought responses by individual tree species are not often correlated with tree species diversity in European forests. *Journal of Applied Ecology*, 53(6), 1725-1734. https://doi.org/10.1111/1365-2664.12745
- Forrester, D. I., & Pretzsch, H. (2015). Tamm review: On the strength of evidence when comparing ecosystem functions of mixtures with monocultures. Forest Ecology and Management, 356, 41–53. https:// doi.org/10.1016/j.foreco.2015.08.016
- Forzieri, G., Girardello, M., Ceccherini, G., Spinoni, J., Feyen, L., Hartmann, H., Beck, P. S. A., Camps-Valls, G., Chirici, G., Mauri, A., & Cescatti, A. (2021). Emergent vulnerability to climate-driven disturbances in

European forests. *Nature Communications*, *12*(1), 1081. https://doi. org/10.1038/s41467-021-21399-7

- Franceschini, T., & Schneider, R. (2014). Influence of shade tolerance and development stage on the allometry of ten temperate tree species. *Oecologia*, 176(3), 739–749. https://doi.org/10.1007/s0044 2-014-3050-3
- Freschet, G. T., Pagès, L., Iversen, C. M., Comas, L. H., Rewald, B., Roumet, C., Klimešová, J., Zadworny, M., Poorter, H., Postma, J. A., Adams, T. S., Bagniewska-Zadworna, A., Bengough, A. G., Blancaflor, E. B., Brunner, I., Cornelissen, J. H. C., Garnier, E., Gessler, A., Hobbie, S. E., ... McCormack, M. L. (2021). A starting guide to root ecology: Strengthening ecological concepts and standardising root classification, sampling, processing and trait measurements. New Phytologist, 232(3), 973-1122. https://doi. org/10.1111/nph.17572
- Gampe, D., Zscheischler, J., Reichstein, M., O'Sullivan, M., Smith, W. K., Sitch, S., & Buermann, W. (2021). Increasing impact of warm droughts on northern ecosystem productivity over recent decades. *Nature Climate Change*, 11(9), 772–779. https://doi.org/10.1038/ s41558-021-01112-8
- Gehring, C. A., Mueller, R. C., & Whitham, T. G. (2006). Environmental and genetic effects on the formation of ectomycorrhizal and arbuscular mycorrhizal associations in cottonwoods. *Oecologia*, 149(1), 158–164. https://doi.org/10.1007/s00442-006-0437-9
- Gillerot, L., Forrester, D. I., Bottero, A., Rigling, A., & Lévesque, M. (2021). Tree Neighbourhood diversity has negligible effects on drought resilience of European Beech, Silver Fir and Norway Spruce. *Ecosystems*, 24(1), 20–36. https://doi.org/10.1007/s10021-020-00501-y
- Grossiord, C. (2020). Having the right neighbors: How tree species diversity modulates drought impacts on forests. New Phytologist, 228(1), 42–49. https://doi.org/10.1111/nph.15667
- Grossiord, C., Granier, A., Ratcliffe, S., Bouriaud, O., Bruelheide, H., Chećko, E., Forrester, D. I., Dawud, S. M., Finér, L., Pollastrini, M., Scherer-Lorenzen, M., Valladares, F., Bonal, D., & Gessler, A. (2014). Tree diversity does not always improve resistance of forest ecosystems to drought. *Proceedings of the National Academy of Sciences of the United States of America*, 111(41), 14812–14815. https://doi.org/ 10.1073/pnas.1411970111
- Haberstroh, S., & Werner, C. (2022). The role of species interactions for forest resilience to drought. *Plant Biology*, 24(7), 1098–1107. https://doi.org/10.1111/plb.13415
- Hari, V., Rakovec, O., Markonis, Y., Hanel, M., & Kumar, R. (2020). Increased future occurrences of the exceptional 2018-2019 Central European drought under global warming. *Scientific Reports*, 10(1), 12207. https://doi.org/10.1038/s41598-020-68872-9
- Hartmann, H., Link, R. M., & Schuldt, B. (2021). A whole-plant perspective of isohydry: Stem-level support for leaf-level plant water regulation. *Tree Physiology*, 41(6), 901–905. https://doi.org/10.1093/ treephys/tpab011
- Hartmann, H., & Trumbore, S. (2016). Understanding the roles of nonstructural carbohydrates in forest trees—From what we can measure to what we want to know. *The New Phytologist*, 211(2), 386– 403. https://doi.org/10.1111/nph.13955
- Heklau, H., Schindler, N., Buscot, F., Eisenhauer, N., Ferlian, O., Prada Salcedo, L. D., & Bruelheide, H. (2021). Mixing tree species associated with arbuscular or ectotrophic mycorrhizae reveals dual mycorrhization and interactive effects on the fungal partners. *Ecology and Evolution*, 11(10), 5424–5440. https://doi.org/10.1002/ece3.7437
- Heklau, H., Schindler, N., Eisenhauer, N., Ferlian, O., & Bruelheide, H. (2023). Temporal variation of mycorrhization rates in a tree diversity experiment. *Ecology and Evolution*, 13(4), 1–14. https://doi.org/ 10.1002/ece3.10002
- Hui, D., Wang, J., Shen, W., Le, X., Ganter, P., & Ren, H. (2014). Near isometric biomass partitioning in forest ecosystems of China. *PLoS One*, 9(1), e86550. https://doi.org/10.1371/journal.pone.0086550

Ingrisch, J., & Bahn, M. (2018). Towards a comparable quantification of resilience. Trends in Ecology & Evolution, 33(4), 251–259. https://doi. org/10.1016/j.tree.2018.01.013

- Isbell, F., Craven, D., Connolly, J., Loreau, M., Schmid, B., Beierkuhnlein, C., Bezemer, T. M., Bonin, C., Bruelheide, H., de Luca, E., Ebeling, A., Griffin, J. N., Guo, Q., Hautier, Y., Hector, A., Jentsch, A., Kreyling, J., Lanta, V., Manning, P., ... Eisenhauer, N. (2015). Biodiversity increases the resistance of ecosystem productivity to climate extremes. *Nature*, 526(7574), 574–577. https://doi.org/10.1038/natur e15374
- Jactel, H., Bauhus, J., Boberg, J., Bonal, D., Castagneyrol, B., Gardiner, B., Gonzalez-Olabarria, J. R., Koricheva, J., Meurisse, N., & Brockerhoff, E. G. (2017). Tree diversity drives forest stand resistance to natural disturbances. Current Forestry Reports, 3(3), 223–243. https://doi. org/10.1007/s40725-017-0064-1
- Jiang, F., Zhu, K., Cadotte, M. W., & Jin, G. (2020). Tree mycorrhizal type mediates the strength of negative density dependence in temperate forests. *Journal of Ecology*, 108(6), 2601–2610. https://doi.org/ 10.1111/1365-2745.13413
- Jourdan, M., Kunstler, G., & Morin, X. (2020). How neighbourhood interactions control the temporal stability and resilience to drought of trees in mountain forests. *Journal of Ecology*, 108(2), 666–677. https://doi.org/10.1111/1365-2745.13294
- Jucker, T., Bouriaud, O., Avacaritei, D., & Coomes, D. A. (2014). Stabilizing effects of diversity on aboveground wood production in forest ecosystems: Linking patterns and processes. *Ecology Letters*, 17(12), 1560–1569. https://doi.org/10.1111/ele.12382
- Jucker, T., Grossiord, C., Bonal, D., Bouriaud, O., Gessler, A., & Coomes, D. A. (2017). Detecting the fingerprint of drought across Europe's forests: Do carbon isotope ratios and stem growth rates tell similar stories? Forest Ecosystems, 4(1), 24. https://doi.org/10.1186/s4066 3-017-0111-1
- Kannenberg, S. A., Novick, K. A., Alexander, M. R., Maxwell, J. T., Moore, D. J. P., Phillips, R. P., & Anderegg, W. R. L. (2019). Linking drought legacy effects across scales: From leaves to tree rings to ecosystems. *Global Change Biology*, 25(9), 2978–2992. https://doi.org/10. 1111/gcb.14710
- Kannenberg, S. A., Schwalm, C. R., & Anderegg, W. R. L. (2020). Ghosts of the past: How drought legacy effects shape forest functioning and carbon cycling. *Ecology Letters*, 23(5), 891–901. https://doi.org/ 10.1111/ele.13485
- Kilpeläinen, J., Barbero-López, A., Vestberg, M., Heiskanen, J., & Lehto, T. (2017). Does severe soil drought have after-effects on arbuscular and ectomycorrhizal root colonisation and plant nutrition? *Plant and Soil*, 418(1–2), 377–386. https://doi.org/10.1007/s1110 4-017-3308-8
- Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). ImerTest package: Tests in linear mixed effects models. *Journal of Statistical Software*, 82(13), 1–26. https://doi.org/10.18637/JSS.V082.113
- Lehto, T., & Zwiazek, J. J. (2011). Ectomycorrhizas and water relations of trees: A review. Mycorrhiza, 21(2), 71–90. https://doi.org/10.1007/ s00572-010-0348-9
- Lenth, R. (2023). *emmeans: Estimated marginal means*. R package Version 1.8.7. https://cran.r-project.org/package=emmeans
- Leuschner, C. (2020). Drought response of European beech (Fagus sylvatica L.)—A review. Perspectives in Plant Ecology, Evolution and Systematics, 47, 125576. https://doi.org/10.1016/j.ppees.2020. 125576
- Liese, R., Lübbe, T., Albers, N. W., & Meier, I. C. (2018). The mycorrhizal type governs root exudation and nitrogen uptake of temperate tree species. *Tree Physiology*, 38(1), 83–95. https://doi.org/10.1093/ treephys/tpx131
- Lloret, F., Keeling, E. G., & Sala, A. (2011). Components of tree resilience: Effects of successive low-growth episodes in old ponderosa pine forests. *Oikos*, 120(12), 1909–1920. https://doi.org/10.1111/j. 1600-0706.2011.19372.x

- Lüdecke, D., Ben-Shachar, M., Patil, I., Waggoner, P., & Makowski, D. (2021). Performance: An R package for assessment, comparison and testing of statistical models. *Journal of Open Source Software*, 6(60), 3139. https://doi.org/10.21105/joss.03139
- Luo, S., Phillips, R. P., Jo, I., Fei, S., Liang, J., Schmid, B., & Eisenhauer, N. (2023). Higher productivity in forests with mixed mycorrhizal strategies. *Nature Communications*, 14(1), 1377. https://doi.org/10. 1038/s41467-023-36888-0
- Luo, S., Schmid, B., De Deyn, G. B., & Yu, S. (2018). Soil microbes promote complementarity effects among co-existing trees through soil nitrogen partitioning. *Functional Ecology*, 32(7), 1879–1889. https:// doi.org/10.1111/1365-2435.13109
- Mao, Z., van der Plas, F., Corrales, A., Anderson-Teixeira, K. J., Bourg, N. A., Chu, C., Hao, Z., Jin, G., Lian, J., Lin, F., Li, B., Luo, W., McShea, W. J., Myers, J. A., Shen, G., Wang, X., Yan, E., Ye, J., Ye, W., ... Wang, X. (2023). Scale-dependent diversity-biomass relationships can be driven by tree mycorrhizal association and soil fertility. *Ecological Monographs*, *93*(2), 1–21. https://doi.org/10.1002/ecm.1568
- Markonis, Y., Kumar, R., Hanel, M., Rakovec, O., Máca, P., & Kouchak, A. A. (2021). The rise of compound warm-season droughts in Europe. *Science Advances*, 7(6), 1–8. https://doi.org/10.1126/sciadv.abb9668
- Messier, C., Bauhus, J., Sousa-Silva, R., Auge, H., Baeten, L., Barsoum, N., Bruelheide, H., Caldwell, B., Cavender-Bares, J., Dhiedt, E., Eisenhauer, N., Ganade, G., Gravel, D., Guillemot, J., Hall, J. S., Hector, A., Hérault, B., Jactel, H., Koricheva, J., ... Zemp, D. C. (2022). For the sake of resilience and multifunctionality, let's diversify planted forests! *Conservation Letters*, 15(1), 1–8. https://doi. org/10.1111/conl.12829
- Midgley, M. G., & Phillips, R. P. (2014). Mycorrhizal associations of dominant trees influence nitrate leaching responses to N deposition. *Biogeochemistry*, 117(2-3), 241–253. https://doi.org/10.1007/ s10533-013-9931-4
- Mohan, J. E., Cowden, C. C., Baas, P., Dawadi, A., Frankson, P. T., Helmick, K., Hughes, E., Khan, S., Lang, A., Machmuller, M., Taylor, M., & Witt, C. A. (2014). Mycorrhizal fungi mediation of terrestrial ecosystem responses to global change: Mini-review. *Fungal Ecology*, 10(1), 3–19. https://doi.org/10.1016/j.funeco.2014.01.005
- Mölder, I., & Leuschner, C. (2014). European beech grows better and is less drought sensitive in mixed than in pure stands: Tree neighbourhood effects on radial increment. *Trees-Structure and Function*, 28(3), 777-792. https://doi.org/10.1007/s00468-014-0991-4
- Morin, X., Fahse, L., de Mazancourt, C., Scherer-Lorenzen, M., & Bugmann, H. (2014). Temporal stability in forest productivity increases with tree diversity due to asynchrony in species dynamics. *Ecology Letters*, 17(12), 1526–1535. https://doi.org/10.1111/ele.12357
- Obladen, N., Dechering, P., Skiadaresis, G., Tegel, W., Keßler, J., Höllerl, S., Kaps, S., Hertel, M., Dulamsuren, C., Seifert, T., Hirsch, M., & Seim, A. (2021). Tree mortality of European beech and Norway spruce induced by 2018–2019 hot droughts in central Germany. *Agricultural and Forest Meteorology*, 307, 108482. https://doi.org/ 10.1016/j.agrformet.2021.108482
- Pardos, M., del Río, M., Pretzsch, H., Jactel, H., Bielak, K., Bravo, F., Brazaitis, G., Defossez, E., Engel, M., Godvod, K., Jacobs, K., Jansone, L., Jansons, A., Morin, X., Nothdurft, A., Oreti, L., Ponette, Q., Pach, M., Riofrío, J., ... Calama, R. (2021). The greater resilience of mixed forests to drought mainly depends on their composition: Analysis along a climate gradient across Europe. *Forest Ecology and Management*, 481, 118687. https://doi.org/10.1016/j.foreco.2020.118687
- Peterson, R. L., & Massicotte, H. B. (2004). Exploring structural definitions of mycorrhizas, with emphasis on nutrient-exchange interfaces. *Canadian Journal of Botany*, 82(8), 1074–1088. https://doi. org/10.1139/b04-071
- Phillips, R. P., Brzostek, E., & Midgley, M. G. (2013). The mycorrhizalassociated nutrient economy: A new framework for predicting carbon-nutrient couplings in temperate forests. *New Phytologist*, 199(1), 41–51. https://doi.org/10.1111/nph.12221

- Pohl, F., Werban, U., Kumar, R., Hildebrandt, A., & Rebmann, C. (2023). Observational evidence of legacy effects of the 2018 drought on a mixed deciduous forest in Germany. *Scientific Reports*, 13(1), 10863. https://doi.org/10.1038/s41598-023-38087-9
- Pretzsch, H. (2020). The course of tree growth. Theory and reality. Forest Ecology and Management, 478, 118508. https://doi.org/10.1016/j. foreco.2020.118508
- Querejeta, J. I., Egerton-Warburton, L. M., & Allen, M. F. (2007). Hydraulic lift may buffer rhizosphere hyphae against the negative effects of severe soil drying in a California oak savanna. *Soil Biology* and Biochemistry, 39(2), 409–417. https://doi.org/10.1016/j.soilbio. 2006.08.008
- Querejeta, J. I., Egerton-Warburton, L. M., & Allen, M. F. (2009). Topographic position modulates the mycorrhizal response of oak trees to interannual rainfall variability. *Ecology*, 90(3), 649–662. https://doi.org/10.1890/07-1696.1
- R Core Team. (2023). R: A language and environment for statistical computing (Version 4.3.1). R Foundation for Statistical Computing. https:// www.r-project.org/
- Rakovec, O., Samaniego, L., Hari, V., Markonis, Y., Moravec, V., Thober, S., Hanel, M., & Kumar, R. (2022). The 2018–2020 multi-year drought sets a new benchmark in Europe. *Earth's Future*, 10(3), 1–11. https:// doi.org/10.1029/2021EF002394
- Ray, T., Delory, B. M., Beugnon, R., Bruelheide, H., Cesarz, S., Eisenhauer, N., Ferlian, O., Quosh, J., von Oheimb, G., & Fichtner, A. (2023). Tree diversity increases productivity through enhancing structural complexity across mycorrhizal types. *Science Advances*, 9(40), eadi2362. https://doi.org/10.1126/sciadv.adi2362
- Read, D. J., & Perez-Moreno, J. (2003). Mycorrhizas and nutrient cycling in ecosystems – A journey towards relevance? *New Phytologist*, 157(3), 475–492. https://doi.org/10.1046/j.1469-8137.2003.00704.x
- Rosling, A., Midgley, M. G., Cheeke, T., Urbina, H., Fransson, P., & Phillips, R. P. (2016). Phosphorus cycling in deciduous forest soil differs between stands dominated by ecto- and arbuscular mycorrhizal trees. New Phytologist, 209(3), 1184–1195. https://doi.org/10.1111/nph.13720
- Sachsenmaier, L., Quosh, J., Ferlian, O., Eisenhauer, N., & Wirth, C. (2024). Data and R code used in: Forest growth resistance and resilience to the 2018–2020 drought depend on tree diversity and mycorrhizal type [dataset]. MyDiv. https://doi.org/10.25829/BPTQ-Z678
- Salomón, R. L., Peters, R. L., Zweifel, R., Sass-Klaassen, U. G. W., Stegehuis, A. I., Smiljanic, M., Poyatos, R., Babst, F., Cienciala, E., Fonti, P., Lerink, B. J. W., Lindner, M., Martinez-Vilalta, J., Mencuccini, M., Nabuurs, G.-J., van der Maaten, E., von Arx, G., Bär, A., Akhmetzyanov, L., ... Steppe, K. (2022). The 2018 European heatwave led to stem dehydration but not to consistent growth reductions in forests. *Nature Communications*, 13(1), 28. https://doi. org/10.1038/s41467-021-27579-9
- Scherer-Lorenzen, M., Potvin, C., Koricheva, J., Schmid, B., Hector, A., Bornik, Z., Reynolds, G., & Schulze, E.-D. (2005). The Design of Experimental Tree Plantations for functional biodiversity research. In Forest diversity and function (pp. 347–376). Springer-Verlag. https://doi.org/10.1007/3-540-26599-6_16
- Schmied, G., Hilmers, T., Mellert, K., Uhl, E., Buness, V., Ambs, D., Steckel, M., Biber, P., Šeho, M., Hoffmann, Y., & Pretzsch, H. (2023). Nutrient regime modulates drought response patterns of three temperate tree species. *Science of the Total Environment*, *868*, 161601. https:// doi.org/10.1016/j.scitotenv.2023.161601
- Schnabel, F., Barry, K. E., Eckhardt, S., Guillemot, J., Geilmann, H., Kahl, A., Moossen, H., Bauhus, J., & Wirth, C. (2024). Neighbourhood species richness and drought-tolerance traits modulate tree growth and δ^{13} C responses to drought. *Plant Biology*, 1-16, 330-345. https://doi.org/10.1111/plb.13611
- Schnabel, F., Liu, X., Kunz, M., Barry, K. E., Bongers, F. J., Bruelheide, H., Fichtner, A., Härdtle, W., Li, S., Pfaff, C., Schmid, B., Schwarz, J. A., Tang, Z., Yang, B., & Bauhus, J. (2021). Species richness stabilizes productivity via asynchrony and drought-tolerance diversity in a

large-scale tree biodiversity experiment. *Science Advances*, 7(51), 11–13. https://doi.org/10.1126/sciadv.abk1643

- Schnabel, F., Purrucker, S., Schmitt, L., Engelmann, R. A., Kahl, A., Richter, R., Seele-Dilbat, C., Skiadaresis, G., & Wirth, C. (2022). Cumulative growth and stress responses to the 2018–2019 drought in a European floodplain forest. *Global Change Biology*, 28(5), 1870– 1883. https://doi.org/10.1111/gcb.16028
- Schnabel, F., Schwarz, J. A., Dănescu, A., Fichtner, A., Nock, C. A., Bauhus, J., & Potvin, C. (2019). Drivers of productivity and its temporal stability in a tropical tree diversity experiment. *Global Change Biology*, 25(12), 4257–4272. https://doi.org/10.1111/gcb.14792
- Schuldt, B., Buras, A., Arend, M., Vitasse, Y., Beierkuhnlein, C., Damm, A., Gharun, M., Grams, T. E. E., Hauck, M., Hajek, P., Hartmann, H., Hiltbrunner, E., Hoch, G., Holloway-Phillips, M., Körner, C., Larysch, E., Lübbe, T., Nelson, D. B., Rammig, A., ... Kahmen, A. (2020). A first assessment of the impact of the extreme 2018 summer drought on Central European forests. *Basic and Applied Ecology*, 45, 86–103. https://doi.org/10.1016/j.baae.2020.04.003
- Schwarz, J., Skiadaresis, G., Kohler, M., Kunz, J., Schnabel, F., Vitali, V., & Bauhus, J. (2020). Quantifying growth responses of trees to drought—A critique of commonly used resilience indices and recommendations for future studies. *Current Forestry Reports*, 6(3), 185-200. https://doi.org/10.1007/s40725-020-00119-2
- Senf, C., Buras, A., Zang, C. S., Rammig, A., & Seidl, R. (2020). Excess forest mortality is consistently linked to drought across Europe. *Nature Communications*, 11(1), 1–8. https://doi.org/10.1038/s4146 7-020-19924-1
- Shovon, T. A., Auge, H., Haase, J., & Nock, C. A. (2024). Positive effects of tree species diversity on productivity switch to negative after severe drought mortality in a temperate forest experiment. *Global Change Biology*, 30(3), 1–14. https://doi.org/10.1111/gcb.17252
- Spinoni, J., Vogt, J. V., Naumann, G., Barbosa, P., & Dosio, A. (2018). Will drought events become more frequent and severe in Europe? International Journal of Climatology, 38(4), 1718–1736. https://doi. org/10.1002/joc.5291
- Tedersoo, L., & Bahram, M. (2019). Mycorrhizal types differ in ecophysiology and alter plant nutrition and soil processes. *Biological Reviews*, 94(5), 1857–1880. https://doi.org/10.1111/brv.12538
- Teste, F. P., Jones, M. D., & Dickie, I. A. (2020). Dual-mycorrhizal plants: Their ecology and relevance. *New Phytologist*, 225(5), 1835–1851. https://doi.org/10.1111/nph.16190
- Thom, D., Buras, A., Heym, M., Klemmt, H.-J., & Wauer, A. (2023). Varying growth response of Central European tree species to the extraordinary drought period of 2018–2020. Agricultural and Forest Meteorology, 338, 109506. https://doi.org/10.1016/j.agrformet. 2023.109506
- Thurm, E. A., Uhl, E., & Pretzsch, H. (2016). Mixture reduces climate sensitivity of Douglas-fir stem growth. Forest Ecology and Management, 376, 205–220. https://doi.org/10.1016/j.foreco.2016.06.020
- Toju, H., Kishida, O., Katayama, N., & Takagi, K. (2016). Networks depicting the fine-scale co-occurrences of fungi in soil horizons. *PLoS One*, 11(11), e0165987. https://doi.org/10.1371/journal.pone.0165987
- Trogisch, S., Liu, X., Rutten, G., Xue, K., Bauhus, J., Brose, U., Bu, W., Cesarz, S., Chesters, D., Connolly, J., Cui, X., Eisenhauer, N., Guo, L., Haider, S., Härdtle, W., Kunz, M., Liu, L., Ma, Z., Neumann, S., ... Bruelheide, H. (2021). The significance of tree-tree interactions for forest ecosystem functioning. *Basic and Applied Ecology*, 55, 33–52. https://doi.org/10.1016/j.baae.2021.02.003
- Vannoppen, A., Treydte, K., Boeckx, P., Kint, V., Ponette, Q., Verheyen, K., & Muys, B. (2020). Tree species diversity improves beech growth and alters its physiological response to drought. *Trees-Structure and*

Function, 34(4), 1059-1073. https://doi.org/10.1007/s00468-020-01981-0

- Vargas, R., Baldocchi, D. D., Querejeta, J. I., Curtis, P. S., Hasselquist, N. J., Janssens, I. A., Allen, M. F., & Montagnani, L. (2010). Ecosystem CO₂ fluxes of arbuscular and ectomycorrhizal dominated vegetation types are differentially influenced by precipitation and temperature. New Phytologist, 185(1), 226–236. https://doi.org/10. 1111/j.1469-8137.2009.03040.x
- Vicente-Serrano, S. M., Beguería, S., & López-Moreno, J. I. (2010). A multiscalar drought index sensitive to global warming: The standardized precipitation evapotranspiration index. *Journal of Climate*, 23(7), 1696–1718. https://doi.org/10.1175/2009JCLI2909.1
- Wagg, C., Jansa, J., Schmid, B., & van der Heijden, M. G. A. (2011). Belowground biodiversity effects of plant symbionts support aboveground productivity. *Ecology Letters*, 14(10), 1001–1009. https://doi.org/10.1111/j.1461-0248.2011.01666.x
- Wagg, C., Jansa, J., Stadler, M., Schmid, B., & van der Heijden, M. G. A. (2011). Mycorrhizal fungal identity and diversity relaxes plant-plant competition. *Ecology*, 92(6), 1303–1313. https://doi.org/10.1890/ 10-1915.1
- Wickham, H. (2016). Data analysis. Springer. https://doi.org/10.1007/ 978-3-319-24277-4_9
- Xu, H., & Zwiazek, J. J. (2020). Fungal aquaporins in ectomycorrhizal root water transport. Frontiers in Plant Science, 11, 1–6. https://doi.org/ 10.3389/fpls.2020.00302
- Zhang, H. Y., Xiao-Tao, L., Hartmann, H., Keller, A., Han, X. G., Trumbore, S., & Phillips, R. P. (2018). Foliar nutrient resorption differs between arbuscular mycorrhizal and ectomycorrhizal trees at local and global scales. *Global Ecology and Biogeography*, 27(7), 875–885. https://doi.org/10.1111/geb.12738
- Zscheischler, J., & Fischer, E. M. (2020). The record-breaking compound hot and dry 2018 growing season in Germany. Weather and Climate Extremes, 29, 100270. https://doi.org/10.1016/j.wace.2020. 100270
- Zscheischler, J., Martius, O., Westra, S., Bevacqua, E., Raymond, C., Horton, R. M., van den Hurk, B., AghaKouchak, A., Jézéquel, A., Mahecha, M. D., Maraun, D., Ramos, A. M., Ridder, N. N., Thiery, W., & Vignotto, E. (2020). A typology of compound weather and climate events. *Nature Reviews Earth and Environment*, 1(7), 333–347. https://doi.org/10.1038/s43017-020-0060-z

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

Appendix S1. Supporting Information on (i) experimental setup, (ii) drought identification, (iii) data cleaning procedure & data preparation and (iv) model supporting information.

How to cite this article: Sachsenmaier, L., Schnabel, F., Dietrich, P., Eisenhauer, N., Ferlian, O., Quosh, J., Richter, R., & Wirth, C. (2024). Forest growth resistance and resilience to the 2018–2020 drought depend on tree diversity and mycorrhizal type. *Journal of Ecology*, 112, 1787–1803. <u>https://</u> doi.org/10.1111/1365-2745.14360